

---

# The Ocean Cannot Absorb Much More CO<sub>2</sub>



By Robert U. Ayres , INSEAD

**Most carbon emissions are absorbed by the ocean, but it's running out of capacity, which could make global temperatures rise even faster.**

Australia's Great Barrier Reef, a 25 million-year-old ecosystem and home to 1,625 species of fish, is on life support. Reports suggest about 93 percent of the reef has succumbed to bleaching, largely the result of climate change.

Spanning 1,400 miles, larger than the United Kingdom, the reef could soon become extinct. Bleaching is the result of increased acidity due to CO<sub>2</sub> entering the oceans from the environment. Currently, it is estimated that 57 percent of new emissions are "dissolved" in the oceans. Moreover, (thanks to warming) the ocean keeps less carbon dioxide in solution. A warmer ocean will dissolve progressively less CO<sub>2</sub>, thus keeping more of the excess CO<sub>2</sub> in the atmosphere. This "carbon-cycle feedback" has different consequences in different general circulation models, ranging from 0.1°C to 1.5°C in increased global temperatures.


As I've pointed out in two [previous articles](#), the true effect of CO2 is fiercely debated and some climate deniers even go so far as to suggest that an increase in emissions is good for the environment, that crop yields will increase as a result of photosynthesis. But ever-improving science tells us that things aren't that simple, especially when it comes to the oceans. So, to finish this series by looking yet again at the evidence, I demonstrate the effect of climate change on the oceans and the implications of this effect.

We know three things about the oceans with certainty:

1. The ocean is warming.
2. The sea level is rising.
3. The ocean is acidifying due to CO2 absorption (which interferes with calcification in organisms, from coral reefs and shellfish to fish bones).

### **The ocean tells the story**

How do we know? First, thanks to the [Argo programme](#), we have increased our coverage of ocean temperatures, from surface to 2,000 meters below surface. With 4,000 floating sensors around the world, we know for certain that the temperature of the oceans is rising as seen below, which confirms historical (albeit more primitive) data.



Second, sea levels have been rising by about 1.7 mm/year since 1901 and about 3.2 mm per year since 1993. The heating of the oceans is accelerating along with CO<sub>2</sub> and temperature increases in the environment. The two main reasons are more water from melting glacier ice on land and from groundwater and thermal expansion due to ocean warming.

Third, we can see ocean acidification in the near death of the Great Barrier Reef and other underwater ecosystems. About 30 percent of the excess carbon dioxide from human activity (fossil fuel combustion) is dissolved in the oceans. It is known that this produces carbonic acid, which subsequently reacts with calcium ions to form bi-carbonate, making it less available to calcification in organisms, such as coral.

### **Taking one for the team**

This is troubling because the ocean has a tendency to reinforce these effects due to its nature. It is important to realise that the oceans are by far the main storage system for heat in the short to medium term, having absorbed 93 percent of the increase in global heating between 1971 and 2010 (IPCC 2014). Oceans can absorb or emit heat much faster than solid rock, and can

store much more (1000 times as much) than the atmosphere. Hence the effective heat storage capacity of the top 700 meters of the oceans, which exhibits measurable seasonal variation, is much larger than the heat storage capacity of either the atmosphere or the land. This is because thermal conductivity of the ground is very low and vertical convection through the crust is almost zero (except during volcanic eruptions), while the mass of the atmosphere is far less than that of the top layer of the ocean. Hence the oceans store much more heat in the summer than the land or atmosphere. This heat is then released during the winter, as warm currents flow toward the poles. There may also be longer cycles, such as El Niños.

The consequences of continuing to let the ocean take one for the team are many. Increased ocean warming alters the “conveyor belt” of surface and deep ocean currents. This could impact the Gulf Stream which warms northern Europe.

Another possible effect of atmospheric warming in the far north would be to thaw some of the “permafrost” area, both above ground and under the Arctic Ocean. Climate warming is happening much faster in the polar regions than in the tropics. (All the climate models show this effect.) The rapid thinning of the ice and likely disappearance (in summer) of the ice in the Arctic Ocean is confirming evidence.

The fact that the poles are getting warmer faster than the tropics means that the north-south temperature gradient is growing smaller and less sharp. That, in turn, is pushing the northern jet stream northward (on average) at the rate of 2 km per year. This would permit both aerobic and anaerobic micro-organism activity under the soil surface to accelerate, releasing both carbon dioxide and methane into the atmosphere. The undersea “cousin” of permafrost on land, methane clathrate, could also start to thaw, releasing methane into the ocean and thence into the atmosphere, resulting in a positive feedback loop.

It seems likely that the rate of heat exchange in the deep oceans (not measured by Argo), depends on the thermohaline (“conveyor belt”) circulation. It is probably much lower than surface heat exchange, since any mixing induced by storms is less important. The rates of mixing vs. conduction and convection are still not well-known. (The residence time of a water molecule in the oceans is estimated to be 3200 years.) An estimated 90 percent of the excess heat warms the oceans, and only 10 percent warms

the land surface. The heat absorbed by the oceans causes thermal expansion of the water. Thermal expansion is one of the three possible causes of global mean sea-level (GMSL) rise.

The other two possible causes of GMSL rise are glacial ice melting and vertical land motion (VLM). Vertical motion results from the removal of the weight of glacial ice in certain terrestrial areas that were once ice-covered, such as northern Canada and Scandinavia. The VLM adjustment is fairly localised. In fact, it is negative (the sea floor is actually sinking, not rising, on average) because the land areas formerly covered by ice are now “springing back” as the weight of glacial ice was removed. The liquid magma under the Earth’s crust gradually rearranges itself as the oceans get heavier and the land gets lighter.

Based on recent evidence, about one third of the GMSL rise – roughly 1.6 mm/year – is due to the thermal expansion of water and two thirds is due to the melting of glacier ice (mainly in Greenland and Antarctica). The quantitative change in ocean mass from glacier ice melting – as opposed to increased volume due to thermal expansion – is now being measured directly, not just estimated from indirect evidence. The ocean mass is now measured from satellites that detect extremely tiny changes in the gravitational attraction over each part of the Earth’s surface. (Gravitational attraction is proportional to mass). This was not possible until after 2002, when ultra-sensitive new instruments for measurement of gravitational force were first utilised in the so-called Gravity Recovery and Climate Experiment (GRACE).

Throughout this series, I have demonstrated that the weight of the evidence is strongly in favour of the theory of anthropogenic climate change, even though there are some weak spots in the theory (water vapour) and the rate of change of sea-levels. The good news is that the temperature rise on the surface of the Earth may be somewhat slower than the worst case scenario. But the bad news is that the positive feedbacks in the system may be significantly greater than heretofore considered, as we can see with the oceans.

The bottom line is that there is no alternative non-anthropogenic theory to explain rising temperatures, melting glaciers, sea level rise and ocean acidification. If we don’t act, the existing mechanisms of the climate will only reinforce the damage already done.

**Find article at**

<https://knowledge.insead.edu/economics-finance/ocean-cannot-absorb-much-more-co2>

---

**About the author(s)**

**Robert U. Ayres** is an Emeritus Professor of Economics and Political Science and Technology Management at INSEAD and the Novartis Chair in Management and the Environment, Emeritus. His most recent book is "[The History and Future of Technology](#)".